arXiv:1509.08741v2 [math.CV] 12 Oct 2016

ON SOME SPECTRAL PROPERTIES OF THE WEIGHTED $\overline{\partial}$-NEUMANN OPERATOR

FRANZ BERGER AND FRIEDRICH HASLINGER

Abstract. We study necessary conditions for compactness of the weighted $\overline{\partial}$-Neumann operator on the space $L^2(\mathbb{C}^n, e^{-\varphi})$ for a plurisubharmonic function φ. Under the assumption that the corresponding weighted Bergman space of entire functions has infinite dimension, a weaker result is obtained by simpler methods. Moreover, we investigate (non-) compactness of the $\overline{\partial}$-Neumann operator for *decoupled* weights, which are of the form $\varphi(z) = \varphi_1(z_1) + \cdots + \varphi_n(z_n)$. More can be said if every $\Delta\varphi_j$ defines a nontrivial doubling measure.

Contents

1. Introduction 1
2. Overview of the results 3
3. A necessary condition for compactness 4
4. A stronger result obtained from Schrödinger operator theory 6
5. Decoupled weights 8
References 10

1. Introduction

We consider the Cauchy-Riemann complex (or Dolbeault complex)

$$0 \to C^\infty(\mathbb{C}^n) \xrightarrow{\overline{\partial}} \Omega^{0,1}(\mathbb{C}^n) \xrightarrow{\overline{\partial}} \Omega^{0,2}(\mathbb{C}^n) \xrightarrow{\overline{\partial}} \cdots \xrightarrow{\overline{\partial}} \Omega^{0,n}(\mathbb{C}^n) \to 0,$$

where $\Omega^{0,q}(\mathbb{C}^n) \coloneqq C^\infty(\mathbb{C}^n, \Lambda^{0,q}T^*(\mathbb{C}^n))$ denotes the space of smooth $(0,q)$-forms on $\mathbb{C}^n$, and the operator $\overline{\partial}$ is defined by

$$\overline{\partial}u \coloneqq \sum_{|J|=q}{}' \sum_{j=1}^n \frac{\partial u_J}{\partial \overline{z}_j}\, d\overline{z}_j \wedge d\overline{z}_J$$

for $u = \sum'_{|J|=q} u_J\, d\overline{z}_J \in \Omega^{0,q}(\mathbb{C}^n)$, and the primed sum indicates that summation is done only over increasing multiindices $J = (j_1, \ldots, j_q)$ of length q, and $d\overline{z}_J \coloneqq d\overline{z}_{j_1} \wedge \cdots \wedge d\overline{z}_{j_q}$.

If $\varphi\colon \mathbb{C}^n \to \mathbb{R}$ is a C^2 function, we denote by $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ the Hilbert space completion of $\Omega^{0,q}_c(\mathbb{C}^n)$ (compactly supported elements of $\Omega^{0,q}(\mathbb{C}^n)$) with respect to the inner product

$$(u,v)_\varphi \coloneqq \sum_{|J|=q}{}' \int_{\mathbb{C}^n} u_J \overline{v_J}\, e^{-\varphi} d\lambda, \tag{1.1}$$

2010 *Mathematics Subject Classification.* Primary 32W05; Secondary 30H20, 35N15, 35P10.
Key words and phrases. weighted $\overline{\partial}$-Neumann problem, decoupled weights, essential spectrum.
Partially supported by the Austrian Science Fund (FWF): P23664.

with λ the Lebesgue measure on $\mathbb{C}^n$. Let $\|\cdot\|_\varphi$ be the associated norm. The $\overline{\partial}$-operator extends maximally to a closed operator on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ by letting it act in the sense of distributions on

$$\mathrm{dom}(\overline{\partial}) \coloneqq \{f \in L^2_{0,q}(\mathbb{C}^n, e^{-\varphi}) : \overline{\partial} f \in L^2_{0,q+1}(\mathbb{C}^n, e^{-\varphi})\}.$$

In other words, $\overline{\partial} f$ is defined to be the distributional derivative of f, viewed as an element of $L^2_{0,q+1}(\mathbb{C}^n, e^{-\varphi})$. Note that the symbol of its formal adjoint with respect to (1.1), which we denote by $\overline{\partial}^t_\varphi$, depends on the weight φ, and we denote the Hilbert space adjoint of $\overline{\partial}$ by $\overline{\partial}^*_\varphi$. For more details on the weighted $\overline{\partial}$-problem, see [12, 11].

The *complex Laplacian* is the Laplacian of the $\overline{\partial}$-complex,

$$\Box^\varphi_{0,q} \coloneqq \overline{\partial}\overline{\partial}^t_\varphi + \overline{\partial}^t_\varphi \overline{\partial} \colon \Omega^{0,q}(\mathbb{C}^n) \to \Omega^{0,q}(\mathbb{C}^n). \tag{1.2}$$

This is an elliptic, formally self-adjoint differential operator, and its *Gaffney extension*, which we also denote by $\Box^\varphi_{0,q}$, is then

$$\Box^\varphi_{0,q} \coloneqq \overline{\partial}\overline{\partial}^*_\varphi + \overline{\partial}^*_\varphi \overline{\partial},$$

understood as an unbounded operator on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ with domain

$$\mathrm{dom}(\Box^\varphi_{0,q}) \coloneqq \{u \in \mathrm{dom}(\overline{\partial}) \cap \mathrm{dom}(\overline{\partial}^*_\varphi) : \overline{\partial} u \in \mathrm{dom}(\overline{\partial}^*_\varphi) \text{ and } \overline{\partial}^*_\varphi u \in \mathrm{dom}(\overline{\partial})\},$$

where $\overline{\partial}^*_\varphi$ is the Hilbert space adjoint of $\overline{\partial}$ for the inner product (1.1), and $\mathrm{dom}(\overline{\partial}^*_\varphi)$ is its domain. It is a positive self-adjoint operator on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ by techniques going back to [9], see also [11]. By completeness of $\mathbb{C}^n$, (1.2) is essentially self-adjoint, so that the Gaffney extension is its only closed symmetric extension, see for instance [11, 16]

The inverse of the restriction of $\Box^\varphi_{0,q}$ to $\mathrm{dom}(\Box^\varphi_{0,q}) \cap \ker(\Box^\varphi_{0,q})^\perp$ is called the *$\overline{\partial}$-Neumann operator* and is denoted by

$$N^\varphi_{0,q} \coloneqq \left(\Box^\varphi_{0,q}|_{\mathrm{dom}(\Box^\varphi_{0,q}) \cap \ker(\Box^\varphi_{0,q})^\perp}\right)^{-1} \colon \mathrm{img}(\Box^\varphi_{0,q}) \to L^2_{0,q}(\mathbb{C}^n, e^{-\varphi}).$$

While $N^\varphi_{0,q}$ is initially only defined on $\mathrm{img}(\Box^\varphi_{0,q})$, we can extend it to a bounded operator on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ if and only if $\Box^\varphi_{0,q}$ has closed range. This is equivalent to $\mathrm{img}(\overline{\partial}) \cap L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ and $\mathrm{img}(\overline{\partial}) \cap L^2_{0,q+1}(\mathbb{C}^n, e^{-\varphi})$ both being closed, and also to the existence of $C > 0$ such that

$$\|u\|^2_\varphi \le C(\|\overline{\partial} u\|^2_\varphi + \|\overline{\partial}^*_\varphi u\|^2_\varphi) \tag{1.3}$$

holds for all $u \in \mathrm{dom}(\overline{\partial}) \cap \mathrm{dom}(\overline{\partial}^*_\varphi) \cap \ker(\Box^\varphi_{0,q})^\perp$, see the arguments in [13, 23, 2]. We are interested in spectral properties of $N^\varphi_{0,q}$, in particular the problem of deciding its compactness in terms of easily accessible properties of φ. We denote by $\sigma(\Box^\varphi_{0,q})$ and $\sigma_e(\Box^\varphi_{0,q})$ the *spectrum* and the *essential spectrum* of $\Box^\varphi_{0,q}$, respectively. The latter set consists of all points in $\sigma(\Box^\varphi_{0,q})$ which are accumulation points of the spectrum or eigenvalues of infinite multiplicity. If $N^\varphi_{0,q}$ is a bounded operator, then it follows from general operator theory that its compactness is equivalent to $\sigma_e(\Box^\varphi_{0,q}) \subseteq \{0\}$. Equivalently, the spectrum of $\Box^\varphi_{0,q}$ may only consist of discrete eigenvalues of finite multiplicity, except possibly allowing for an infinite dimensional kernel.

One important property of the $\overline{\partial}$-Neumann operator lies in the fact that it can be used to compute the norm-minimal solution to the inhomogeneous equation $\overline{\partial} u = f$, for a given $\overline{\partial}$-closed $(0, q+1)$-form f. Indeed, this solution is then given by $u = \overline{\partial}^*_\varphi N_{0,q+1} f$, and many operator theoretic properties such as compactness transfer from N to the minimal solution operator $\overline{\partial}^*_\varphi N$. We refer the reader to the literature, for example [23, 11].

An important tool is the (weighted) *Kohn-Morrey-Hörmander formula*,

$$\|\bar\partial u\|_\varphi^2 + \|\bar\partial_\varphi^* u\|_\varphi^2 = \sum_{|J|=q}{}' \sum_{j=1}^n \left\| \frac{\partial u_J}{\partial \overline{z}_j} \right\|_\varphi^2 + \sum_{|K|=q-1}{}' \sum_{j,k=1}^n \int_{\mathbb{C}^n} \frac{\partial^2 \varphi}{\partial z_j \partial \overline{z}_k}\, u_{jK} \overline{u_{kK}}\, e^{-\varphi} d\lambda, \tag{1.4}$$

see [13], [11, p. 109], or [23, Proposition 2.4], which is valid a priori for $u \in \Omega_c^{0,q}(\mathbb{C}^n)$ with $1 \le q \le n$, and by extension also for $u \in \mathrm{dom}(\bar\partial) \cap \mathrm{dom}(\bar\partial_\varphi^*)$, since $\Omega_c^{0,q}(\mathbb{C}^n)$ is dense in the latter space for the norm $u \mapsto (\|u\|_\varphi^2 + \|\bar\partial u\|_\varphi^2 + \|\bar\partial_\varphi^* u\|_\varphi^2)^{1/2}$. Again, the reason for this is that $\mathbb{C}^n$ is complete for the Euclidean metric.

Remark 1.1. (i) From (1.4), one derives a sufficient condition for boundedness of $N_{0,q}^\varphi$, which is that φ be plurisubharmonic, and

$$\liminf_{z\to\infty} s_q(z) > 0,$$

where $s_q(z)$ is the sum of the q smallest eigenvalues of the complex Hessian

$$M_\varphi(z) := \left(\frac{\partial^2 \varphi}{\partial z_j \partial \overline{z}_k}(z) \right)_{j,k=1}^n$$

of φ. W refer to [11] for the details. From this condition it also follows that $\Box_{0,q}^\varphi$ is injective, so that $N_{0,q}^\varphi$ is the bounded inverse of $\Box_{0,q}^\varphi$.[1] (ii) A sufficient condition for $N_{0,q}^\varphi$ being compact is the following: Suppose that $\varphi \in C^2(\mathbb{C}^n, \mathbb{R})$ and that the sum $s_q(z)$ of the smallest q eigenvalues of $M_\varphi(z)$ has the property

$$\lim_{z\to\infty} s_q(z) = +\infty. \tag{1.5}$$

Then $N_{0,q}^\varphi$ is compact. Indeed, (1.4) together with some linear algebra implies that $\|\bar\partial u\|_\varphi^2 + \|\bar\partial_\varphi^* u\|_\varphi^2 \ge (s_q u, v)_\varphi$ for all $u, v \in \mathrm{dom}(\bar\partial) \cap \mathrm{dom}(\bar\partial_\varphi^*) \cap L_{0,q}^2(\mathbb{C}^n, e^{-\varphi})$, and hence the diverging of s_q implies compactness, see [10, 11] or also [20] for the details. Further recent results concerning compactness of the $\bar\partial$-Neumann operator for the weighted problem can be found in [6, 7].

(iii) Boundedness and compactness of the $\bar\partial$-Neumann operator "percolate" up the $\bar\partial$-complex: if $N_{0,q}^\varphi$ is bounded or compact, then $N_{0,q+1}^\varphi$ has the same property, see [11].

2. Overview of the results

In this article we will first derive a necessary condition for compactness to hold, which is weaker than (1.5): Under the assumption that the *weighted Bergman space*

$$A^2(\mathbb{C}^n, e^{-\varphi}) := \left\{ f \colon \mathbb{C}^n \to \mathbb{C} : f \text{ entire and } \int_{\mathbb{C}^n} |f|^2\, e^{-\varphi} d\lambda < \infty \right\}$$

has infinite dimension (see Remark 3.2 for a known sufficient condition for this) and $N_{0,q}^\varphi$ is compact for some $1 \le q \le n$, then

$$\limsup_{z\to\infty} \mathrm{tr}(M_\varphi(z)) = +\infty, \tag{2.1}$$

[1] Actually, for the Laplacian to be injective it suffices to have φ plurisubharmonic and $M_\varphi(z) \neq 0$ for some $z \in \mathbb{C}^n$, compare with the arguments in the proof of Theorem 5.1.

or, in other words, $\operatorname{tr}(M_\varphi)$ is unbounded. A stronger result, obtained with different methods, is presented in section 4: it says that if $N^\varphi_{0,q}$ is compact (regardless of dimensionality of the Bergman space), then

$$\lim_{z\to\infty} \int_{B_1(z)} \operatorname{tr}(M_\varphi)^2 \, d\lambda = +\infty.$$

Of course, this also implies (2.1).

In section 5 we will consider weights which are *decoupled*, i.e., of the form

$$\varphi(z_1, \dots, z_n) = \varphi_1(z_1) + \cdots + \varphi_n(z_n).$$

Decoupled weights are known from [12] to be an obstruction to compactness of the $\overline\partial$-Neumann operator on $(0,1)$-forms, provided at least one φ_j gives rise to an infinite dimensional weighted Bergman space (of entire functions on $\mathbb{C}$). By using results of [2, 17, 18], we will characterize compactness of $N^\varphi_{0,q}$ for this class of weights, under the additional assumption that each $\varphi_j : \mathbb{C} \to \mathbb{R}$ is a subharmonic function such that $\Delta\varphi_j$ defines a nontrivial doubling measure. This means that φ_j is not harmonic and there is $C > 0$ such that

$$\int_{B_{2r}(z)} \Delta\varphi_j \, d\lambda \le C \int_{B_r(z)} \Delta\varphi_j \, d\lambda$$

for all $z \in \mathbb{C}$ and $r > 0$. As an example, $\varphi(z) := |z|^\alpha$ for $\alpha > 0$ satisfies these conditions. It turns out that $N^\varphi_{0,q}$ is never compact for $0 \le q \le n-1$, and compactness of $N^\varphi_{0,n}$ depends on the asymptotics of the integral $\int_{B_1(z)} \operatorname{tr}(M_\varphi)\, d\lambda$ as $z \to \infty$. This will also give examples of weights φ such that (2.1) is not sufficient for compactness of N^φ.

3. A necessary condition for compactness

To derive a necessary condition for compactness of $N^\varphi_{0,q}$, we will apply $\Box^\varphi_{0,q}$ to $(0,q)$-forms with coefficients (for the standard basis) belonging to $A^2(\mathbb{C}^n, e^{-\varphi})$. The point is that $\Box^\varphi_{0,q}$ restricts to a multiplication operator on this space.

Theorem 3.1. *Let $\varphi\colon \mathbb{C}^n \to \mathbb{R}$ be a plurisubharmonic C^2 function and suppose that the corresponding weighted space $A^2(\mathbb{C}^n, e^{-\varphi})$ of entire function is infinite dimensional. If there is $1 \le q \le n$ such that the $\overline\partial$-Neumann operator $N^\varphi_{0,q}$ is compact, then*

$$\limsup_{z\to\infty} \operatorname{tr}(M_\varphi(z)) = +\infty. \tag{3.1}$$

Proof. Let g be a smooth $(0,q)$-form, $g = \sum'_{|J|=q} g_J \, d\overline z_J \in \Omega^{0,q}(\mathbb{C}^n)$. If the coefficients g_J of g are holomorphic, then $g \in \ker(\overline\partial\colon \Omega^{0,q}(\mathbb{C}^n) \to \Omega^{0,q+1}(\mathbb{C}^n))$, and $\Box^\varphi_{0,q} g$ as in (1.2) is given by

$$\begin{aligned}
\overline\partial\overline\partial^t_\varphi g &= \overline\partial\Big(- \sum_{|K|=q-1}{}' \sum_{k=1}^n \Big(\frac{\partial}{\partial z_k} - \frac{\partial\varphi}{\partial z_k}\Big) g_{kK} \, d\overline z_K \Big) \\
&= -\sum_{j=1}^n \sum_{|K|=q-1}{}' \sum_{k=1}^n \Big(\frac{\partial^2}{\partial\overline z_j \partial z_k} - \frac{\partial\varphi}{\partial z_k}\frac{\partial}{\partial\overline z_j} - \frac{\partial^2\varphi}{\partial\overline z_j\partial z_k}\Big) g_{kK} \, d\overline z_j \wedge d\overline z_K,
\end{aligned}$$

which reduces to

$$\sum_{|K|=q-1}{}' \sum_{j,k=1}^n \frac{\partial^2\varphi}{\partial\overline z_j \partial z_k} g_{kK} \, d\overline z_j \wedge d\overline z_K, \tag{3.2}$$

see [11] for the computation of $\bar\partial^t_\varphi$. If $\widetilde{\Box}^\varphi_{0,q}$ denotes the maximal closed extension of (1.2) to an operator on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$, defined in the sense of distributions on

$$\mathrm{dom}(\widetilde{\Box}^\varphi_{0,q}) := \{u \in L^2_{0,q}(\mathbb{C}^n, e^{-\varphi}) : \Box^\varphi_{0,q} u \in L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})\},$$

then $\widetilde{\Box}^\varphi_{0,q}$ is symmetric and hence must equal the Gaffney extension.

Now assume that (3.1) is wrong, so that there exists $C \geq 0$ such that

$$\mathrm{tr}(M_\varphi(z)) \leq C \tag{3.3}$$

for all $z \in \mathbb{C}^n$. Since φ is assumed to be plurisubharmonic, M_φ is nonnegative and hence (3.3) is equivalent to all entries of M_φ being bounded. Indeed, $\mathrm{tr}(M^*_\varphi M_\varphi) = \mathrm{tr}(M^2_\varphi) \leq \mathrm{tr}(M_\varphi)^2$, and the Hilbert-Schmidt norm $A \mapsto \mathrm{tr}(A^*A)^{1/2}$ on the space of complex $n \times n$ matrices is equivalent to the max norm (i.e., the maximum of the entries). It now follows from (3.2) that all $(0,q)$-forms g with coefficients in $A^2(\mathbb{C}^n, e^{-\varphi})$ belong to $\mathrm{dom}(\widetilde{\Box}^\varphi_{0,q}) = \mathrm{dom}(\Box^\varphi_{0,q})$, and

$$\Box^\varphi_{0,q} g = {\sum_{|K|=q-1}}' \sum_{j,k=1}^n \frac{\partial^2 \varphi}{\partial \overline{z}_j \partial z_k} g_{kK}\, d\overline{z}_j \wedge d\overline{z}_K.$$

Hence, if all such g belong to a bounded set of $L^2_{(0,q)}(\mathbb{C}^n, e^{-\varphi})$, the image under $\Box^\varphi_{0,q}$ of this bounded set will be again bounded in $L^2_{(0,q)}(\mathbb{C}^n, e^{-\varphi})$. In addition we have

$$N^\varphi_{0,q}\left({\sum_{|K|=q-1}}' \sum_{j,k=1}^n \frac{\partial^2 \varphi}{\partial \overline{z}_j \partial z_k} g_{kK}\, d\overline{z}_j \wedge d\overline{z}_K \right) = g,$$

so that the identity on the (infinite dimensional) space of $(0,q)$-forms with coefficients in $A^2(\mathbb{C}^n, e^{-\varphi})$ is the composition of the compact operator $N^\varphi_{0,q}$ and a bounded operator, hence it is compact. This contradicts $\dim(A^2(\mathbb{C}^n, e^{-\varphi})) = \infty$. □

Remark 3.2. (i) To obtain Theorem 3.1, one could also use the fact that compactness of $N^\varphi_{0,q}$ "percolates up" the weighted $\bar\partial$-complex, see [11]: if $1 \leq q \leq n-1$ and $N^{0,q}_\varphi$ is compact, then $N^{0,q+1}_\varphi$ is also compact. At the upper end of the $\bar\partial$-complex, one has

$$\begin{aligned} \Box^\varphi_{0,n} g &= \bar\partial \bar\partial^t_\varphi g \\ &= \bar\partial\Big(- {\sum_{|K|=n-1}}' \sum_{k=1}^n \Big(\frac{\partial}{\partial z_k} - \frac{\partial \varphi}{\partial z_k} \Big) g_{kK}\, d\overline{z}_K \Big) \\ &= -\frac{1}{4}\Delta g + \sum_{j=1}^n \frac{\partial \varphi}{\partial z_j} \frac{\partial g}{\partial \overline{z}_j} + \frac{1}{4}(\Delta\varphi) g, \end{aligned} \tag{3.4}$$

for $g \in \Omega^{0,n}(\mathbb{C}^n)$, where we identify the $(0,n)$-form

$$g = \widetilde{g}\, d\overline{z}_1 \wedge \cdots \wedge d\overline{z}_n \in \Omega^{0,n}(\mathbb{C}^n)$$

with its coefficient $\widetilde{g} \in C^\infty(\mathbb{C}^n)$. If we consider $(0,n)$-forms g with coefficient belonging to $A^2(\mathbb{C}^n, e^{-\varphi})$, then $g \in \Omega^{0,n}(\mathbb{C}^n) \cap L^2_{0,n}(\mathbb{C}^n, e^{-\varphi})$ and, by (3.4), we have

$$\Box^\varphi_{0,n} g = \frac{1}{4}(\Delta\varphi) g = \mathrm{tr}(M_\varphi) g,$$

and one can now use the same argument as before to reach a contradiction.

(ii) Let $n = 1$ and suppose that $\varphi\colon \mathbb{C} \to \mathbb{R}$ is a subharmonic C^2 function such that the Bergman space $A^2(\mathbb{C}, e^{-\varphi})$ is infinite dimensional. The complex Laplacian on $(0,1)$-forms equals $\Box^{\varphi}_{0,1} = \overline{\partial}\overline{\partial}^*_{\varphi}$, and the necessary condition (3.1) for compactness of the $\overline{\partial}$-Neumann operator becomes

$$\limsup_{z\to\infty} \Delta\varphi(z) = +\infty. \tag{3.5}$$

If one supposes that $\Delta\varphi$ defines a doubling measure, the condition

$$\lim_{z\to\infty} \int_{B_1(z)} \Delta\varphi \, d\lambda = +\infty \tag{3.6}$$

is known to be both necessary and sufficient for compactness of the $\overline{\partial}$-Neumann operator, see [12] and [18]. Of course, (3.6) implies (3.5), as it should.

(iii) A sufficient condition for the Bergman space $A^2(\mathbb{C}^n, e^{-\varphi})$ to have infinite dimension is given in [21, Lemma 3.4]: If $\varphi\colon \mathbb{C}^n \to \mathbb{R}$ is a smooth function such that

$$\lim_{z\to\infty} |z|^2 s_1(z) = +\infty, \tag{3.7}$$

where s_1 is the smallest eigenvalue of M_φ, then $A^2(\mathbb{C}^n, e^{-\varphi})$ has infinite dimension. Condition (3.7) is not sharp: the function

$$\varphi(z_1, z_2) = |z_1|^4 + |z_1 z_2|^2$$

on $\mathbb{C}^2$ does not satisfy (3.7). Nevertheless, the corresponding space $A^2(\mathbb{C}^2, e^{-\varphi})$ is of infinite dimension since it contains all polynomials in z_1, see the computation in [11, p. 125].

4. A stronger result obtained from Schrödinger operator theory

Let $u = u\, d\overline{z}_1 \wedge \cdots \wedge d\overline{z}_n$ be a smooth $(0,n)$-form belonging to the domain of $\Box^{\varphi}_{0,n}$. For $1 \le j \le n$ denote by K_j the increasing multiindex $K_j := (1, \ldots, j-1, j+1, \ldots, n)$ of length $n-1$. Then

$$\overline{\partial}^t_{\varphi} u = \sum_{j=1}^{n} (-1)^{j+1} \left(\frac{\partial\varphi}{\partial z_j} u - \frac{\partial u}{\partial z_j} \right) d\overline{z}_{K_j}.$$

Hence

$$\begin{aligned} \overline{\partial}\overline{\partial}^t_{\varphi} u &= \left[\sum_{j=1}^{n} \frac{\partial}{\partial \overline{z}_j} \left(\frac{\partial\varphi}{\partial z_j} u - \frac{\partial u}{\partial z_j} \right) \right] d\overline{z}_1 \wedge \cdots \wedge d\overline{z}_n \\ &= \left[\sum_{j=1}^{n} \left(\frac{\partial^2\varphi}{\partial z_j \partial \overline{z}_j} u + \frac{\partial\varphi}{\partial z_j} \frac{\partial u}{\partial \overline{z}_j} - \frac{\partial^2 u}{\partial z_j \partial \overline{z}_j} \right) \right] d\overline{z}_1 \wedge \cdots \wedge d\overline{z}_n. \end{aligned}$$

Conjugation with the unitary operator $L^2(\mathbb{C}^n, e^{-\varphi}) \to L^2(\mathbb{C}^n)$ of multiplication by $e^{-\varphi/2}$ gives

$$e^{-\varphi/2} \Box^{\varphi}_{0,n} e^{\varphi/2} f = \sum_{j=1}^{n} \Big(- \frac{\partial^2 f}{\partial z_j \partial \overline{z}_j} - \frac{1}{2} \frac{\partial\varphi}{\partial \overline{z}_j} \frac{\partial f}{\partial z_j} + \frac{1}{2} \frac{\partial\varphi}{\partial z_j} \frac{\partial f}{\partial \overline{z}_j} + \frac{1}{4} \frac{\partial\varphi}{\partial \overline{z}_j} \frac{\partial\varphi}{\partial z_j} f + \frac{1}{2} \frac{\partial^2\varphi}{\partial z_j \partial \overline{z}_j} f \Big), \tag{4.1}$$

where $f \in L^2(\mathbb{C}^n)$ and we just wrote down the coefficient of the corresponding $(0,n)$-form. This operator can be expressed by real variables in the form

$$e^{-\varphi/2} \Box^{\varphi}_{0,n} e^{\varphi/2} f = \frac{1}{4} (-\Delta_A + V) f, \tag{4.2}$$

with

$$\Delta_A := \sum_{j=1}^{n} \Big[\Big(-\frac{\partial}{\partial x_j} - \frac{i}{2}\frac{\partial \varphi}{\partial y_j} \Big)^2 + \Big(-\frac{\partial}{\partial y_j} + \frac{i}{2}\frac{\partial \varphi}{\partial x_j} \Big)^2 \Big],$$

and $V := 2\operatorname{tr}(M_\varphi)$. It follows that $-\Delta_A + V$ is a Schrödinger operator on $L^2(\mathbb{R}^{2n})$ with magnetic vector potential

$$A := \frac{1}{2}\Big(-\frac{\partial \varphi}{\partial y_1}, \frac{\partial \varphi}{\partial x_1}, \dots, -\frac{\partial \varphi}{\partial y_n}, \frac{\partial \varphi}{\partial x_n} \Big),$$

where $z_j = x_j + iy_j$, $j = 1, \dots, n$, and non-negative electric potential V in the case where φ is plurisubharmonic. Note that A corresponds to the real 1-form $-\frac{i}{2}(\partial - \bar\partial)\varphi$, and the associated magnetic field is $dA = i\partial\bar\partial\varphi$. By essential self-adjointness, the Gaffney extension of $\Box^\varphi_{0,n}$ agrees with the unique self-adjoint extension of the Schrödinger operator. We can now apply a result of Iwatsuka [15] to obtain:

Theorem 4.1. *Let $\varphi\colon \mathbb{C}^n \to \mathbb{R}$ be smooth and plurisubharmonic. Then $\Box^\varphi_{0,n}$ has compact resolvent on $L^2_{0,n}(\mathbb{C}^n, e^{-\varphi})$ if and only if the magnetic Schrödinger operator $-\Delta_A + V$ has compact resolvent on $L^2(\mathbb{R}^{2n})$. In particular, if there is $1 \le q \le n$ such that $\Box^\varphi_{0,q}$ has compact resolvent, then*

$$\lim_{z\to\infty} \int_{B(z,1)} \operatorname{tr}(M_\varphi)^2 \, d\lambda = +\infty \tag{4.3}$$

Proof. It remains to prove the last assertion. If $\Box^\varphi_{0,q}$ has compact resolvent, then so does $\Box^\varphi_{0,n}$, see Remark 1.1, and if the magnetic Schrödinger operator has compact resolvent, then

$$\int_{B(z,1)} (|M_\varphi|^2 + 2\operatorname{tr}(M_\varphi)) d\lambda \to \infty \tag{4.4}$$

as $z \to \infty$ by [15, Theorem 5.2], where $|B|^2 := \sum_{j,k=1}^n |B_{j,k}|^2$, since the magnetic field is $i\partial\bar\partial\varphi$. Because $M_\varphi \ge 0$, the above is equivalent to $\int_{B(z,1)} \big(\operatorname{tr}(M_\varphi)^2 + \operatorname{tr}(M_\varphi)\big)\, d\lambda \to \infty$ as $z \to \infty$. By Hölder's inequality,

$$\int_{B(z,1)} \operatorname{tr}(M_\varphi)\, d\lambda \le |B(z,1)|^{1/2} \Big(\int_{B(z,1)} \operatorname{tr}(M_\varphi)^2 d\lambda \Big)^{1/2},$$

and we see that (4.4) implies $f\big(\int_{B(z,1)} \operatorname{tr}(M_\varphi)^2\, d\lambda\big) \to \infty$ as $z \to \infty$, where $f\colon [0,\infty) \to [0,\infty)$ is $f(t) := t + t^{1/2}$. Because f is a monotone bijection, this is equivalent to $\int_{B(z,1)} \operatorname{tr}(M_\varphi)^2\, d\lambda \to \infty$ as $z \to \infty$. □

Remark 4.2. (i) The above condition (4.3) is not a sufficient condition for compactness of $N^\varphi_{0,q}$ in general, see Remark 5.2 below.

(ii) If $\varphi\colon \mathbb{C}^n \to \mathbb{R}$ is such that $\operatorname{tr}(M_\varphi)$ satisfies the reverse Hölder condition,

$$\Big(\fint_B \operatorname{tr}(M_\varphi)^r \, d\lambda \Big)^{1/r} \le C \fint_B \operatorname{tr}(M_\varphi)\, d\lambda$$

for some $r \ge 2$, some $C > 0$, and all balls $B \subseteq \mathbb{C}^n$, where $\fint_B$ denotes the average over B for the Lebesgue measure, then Hölder's inequality implies that (4.3) can be replaced by the formally weaker condition of

$$\lim_{z\to\infty} \int_{B(z,1)} \operatorname{tr}(M_\varphi)\, d\lambda = +\infty. \tag{4.5}$$

The class of functions satisfying one of these reverse Hölder conditions equals $A_\infty := \bigcup_{p\geq 1} A_p$, where A_p are the *Muckenhoupt classes*, see [22, Theorem 3, p. 212]. Every positive polynomial belongs to A_∞. In fact, $|P|^a \in A_p$ for $p > 1$ if $-1 < ad < p-1$, where d is the degree of P, see [22, 6.5, p 219].

5. Decoupled weights

We will now consider weight functions φ which are *decoupled*, meaning that

$$\varphi(z) = \varphi_1(z_1) + \cdots + \varphi_n(z_n) \tag{5.1}$$

for functions $\varphi_j \colon \mathbb{C} \to \mathbb{R}$. Then

$$\sigma(\Box^\varphi_{0,q}) = \bigcup_{\substack{q_1+\cdots+q_n=q \\ q_j\in\{0,1\}}} (\sigma(\Box^{\varphi_1}_{0,q_1}) + \cdots + \sigma(\Box^{\varphi_n}_{0,q_n})) \tag{5.2}$$

and

$$\sigma_e(\Box^\varphi_{0,q}) = \bigcup_{\substack{q_1+\cdots+q_n=q \\ q_j\in\{0,1\}}} \bigcup_{j=1}^{n} \Big(\sigma_e(\Box^{\varphi_j}_{0,q_j}) + \sum_{k\neq j} \sigma(\Box^{\varphi_k}_{0,q_k})\Big). \tag{5.3}$$

Here, $\Box^{\varphi_j}_{0,q_j}$ denotes the complex Laplacian for the weight φ_j (of one variable) on $\mathbb{C}$. This can be seen as a special case of [2, Theorem 5.5]. Indeed, the $\overline\partial$-operator acting on $L^2_{0,q}(\mathbb{C}^n, e^{-\varphi})$ can be understood geometrically as the $\overline\partial^E$-operator for the trivial line bundle $E := \mathbb{C}^n \times \mathbb{C} \to \mathbb{C}^n$, and with Hermitian metric $\langle (z,v_1),(z,v_2)\rangle := v_1\overline{v_2}\, e^{-\varphi(z)}$ on E. If now E_j is $\mathbb{C}\times\mathbb{C}\to\mathbb{C}$ with metric given by $\varphi_j\colon \mathbb{C}\to\mathbb{R}$, then it is easy to see that under the isomorphism

$$(\mathbb{C}^n \times \mathbb{C} \to \mathbb{C}) \cong \pi_1^* E_1 \otimes \cdots \otimes \pi_n^* E_n$$

the trivial line bundle on $\mathbb{C}^n$ carries the metric given by (5.1), where $\pi_j\colon \mathbb{C}^n \to \mathbb{C}$ is the projection onto the jth factor. For the details on the general definition of $\overline\partial^E$ and its properties, we refer to [24, 16, 14, 8]. We note that equation (5.2) has also appeared in [4].

In the following Theorem 5.1, we will consider the case where all $\Delta\varphi_j$ define nontrivial doubling measures. It is known from [18] (or [12, Theorem 2.3], with slightly stronger assumptions) that, under these conditions, $N^{\varphi_j}_{0,1}$ (for the one-variable weights) is compact if and only if

$$\lim_{z\to\infty} \int_{B_1(z)} \Delta\varphi_j \, d\lambda = +\infty. \tag{5.4}$$

holds. Using this condition and the above results, we can characterize compactness of the $\overline\partial$-Neumann operator in terms of the decoupled weight:

Theorem 5.1. *Let $\varphi_j \in C^2(\mathbb{C},\mathbb{R})$ for $1 \leq j \leq n$ with $n \geq 2$, and set $\varphi(z_1,\dots,z_n) := \varphi_1(z_1) + \cdots + \varphi_n(z_n)$. Assume that all φ_j are subharmonic and such that $\Delta\varphi_j$ defines a nontrivial doubling measure, then*

(i) $\dim(\ker(\Box^\varphi_{0,0})) = \dim(A^2(\mathbb{C}^n, e^{-\varphi})) = \infty$,
(ii) $\ker(\Box^\varphi_{0,q}) = 0$ *for* $q \geq 1$,
(iii) $N^\varphi_{0,q}$ *is bounded for* $0 \leq q \leq n$,
(iv) $N^\varphi_{0,q}$ *with* $0 \leq q \leq n-1$ *is not compact, and*

(v) $N^{\varphi}_{0,n}$ is compact if and only if

$$\lim_{z\to\infty}\int_{B_1(z)}\operatorname{tr}(M_\varphi)\,d\lambda=\infty. \tag{5.5}$$

Proof. From [17, Theorem C], it follows from our assumptions on φ that $\bar\partial$ has closed range in $L^2_{0,1}(\mathbb{C},e^{-\varphi_j})$ for all j. From this it also follows that $\bar\partial$ has closed range in $L^2_{0,q}(\mathbb{C}^n,e^{-\varphi})$ for all $0\le q\le n$, see [5, Theorem 4.5] or [3, Corollary 2.15], which implies that the $\bar\partial$-Neumann operator is bounded.

Moreover, $\ker(\Box^{\varphi_j}_{0,1})=0$ for all j. In fact, in this simple case, we obtain from the Kohn-Morrey-Hörmander formula (1.4)

$$\|\Box^{\varphi_j}_{0,1}u\|^2=\|\bar\partial^*_{\varphi_j}u\|^2\ge\frac14\int_{\mathbb{C}}\Delta\varphi_j|u_1|^2\,e^{-\varphi_j}d\lambda \tag{5.6}$$

for all forms $u=u_1d\bar z\in\operatorname{dom}(\bar\partial^*_{\varphi_j})$. If $u\in\ker(\Box^{\varphi_j}_{0,1})$, then u is smooth by elliptic regularity, and if z_0 is such that $\Delta\varphi_j(z_0)>0$ (note that $\Delta\varphi_j\ge0$ everywhere by subharmonicity), then $u=0$ in a neighborhood of z_0. But then $u=0$ everywhere since $\mathbb{C}$ is connected and by using a unique continuation principle of Aronszajn, see [1] or [8, p. 333]. We also have $\ker(\Box^{\varphi}_{0,q})=0$ for all $q\ge1$, either by combining the above with the *Künneth formula*,

$$\ker(\Box^{\varphi}_{0,q})\cong\bigoplus_{q_1+\cdots+q_n=q}\ker(\Box^{\varphi_1}_{0,q_1})\,\hat\otimes\cdots\hat\otimes\,\ker(\Box^{\varphi_n}_{0,q_n}), \tag{5.7}$$

see [3, Corollary 2.15] or [5, Theorem 4.5], where $\hat\otimes$ denotes the Hilbert space tensor product, or directly by using the same argument as above, i.e., applying (1.4) to the higher dimensional problem and using that $M_\varphi\neq0$ at one point.

As a nontrivial doubling measure, $\Delta\varphi_j$ satisfies $\int_{\mathbb{C}}\Delta\varphi_j\,d\lambda=\infty$. Consequently, the weighted Bergman space $A^2(\mathbb{C},e^{-\varphi_j})$ has infinite dimension by [19, Theorem 3.2]. This also implies, by (5.7), that $\dim(\ker(\Box^{\varphi}_{0,0}))=\infty$. On the other hand, $\dim(A^2(\mathbb{C},e^{-\varphi_j}))=\infty$ for at least one $1\le j\le n$ implies that $N^{\varphi}_{0,q}$ cannot be compact for $0\le q\le n-1$, see [2, Theorem 5.6]: indeed, $0\in\sigma_e(\Box^{\varphi_j}_{0,0})$ and hence (5.3) implies that the infinite sets $\sigma(\Box^{\varphi_k}_{0,0})$ for $j\neq k$ are contained in $\sigma_e(\Box^{\varphi}_{0,q})$. This finishes the proof of (i) to (iv). Again by [2, Theorem 5.6], $N^{\varphi}_{0,n}$ is compact if and only if all $N^{\varphi_j}_{0,1}$ are compact, which is the case if and only if (5.4) holds for all $1\le j\le n$. It remains to show that this is equivalent to (5.5). By a simple scaling argument, the claim is equivalent to

$$\int_{B_1(z_1)\times\cdots\times B_n(z_n)}\operatorname{tr}(M_\varphi)\,d\lambda=\frac{\pi^{n-1}}{4}\sum_{j=1}^n\int_{B_1(z_j)}\Delta\varphi_j\,d\lambda\to\infty\quad\text{as}\quad z=(z_1,\dots,z_n)\to\infty, \tag{5.8}$$

and if (5.4) holds for all $1\le j\le n$, then (5.8) is also satisfied. Conversely, if (5.8) is true, then choosing $z=\zeta e_k$ with $\zeta\in\mathbb{C}$ and e_k the kth standard basis vector of $\mathbb{C}^n$ implies

$$\int_{B_1(\zeta)}\Delta\varphi_k\,d\lambda+\sum_{j\neq k}\int_{B_1(0)}\Delta\varphi_j\,d\lambda\to\infty\quad\text{as}\quad\zeta\to\infty,$$

so that $\lim_{\zeta\to\infty}\int_{B_1(\zeta)}\Delta\varphi_k\,d\lambda=\infty$ since the second term is bounded. This shows (v) and concludes the proof. □

Remark 5.2. (i) The doubling condition is satisfied if the $\Delta\varphi_j$ belong to A_∞, see [22, p. 196], where we recall from Remark 4.2 that A_∞ is the union of the Muckenhoupt classes. As an example, $z\mapsto|z|^\alpha$ is in A_p for $p>1$ if and only if $-2<\alpha<2(p-1)$, and defines a doubling measure

for $-2 < \alpha$, cf. [22, 6.4, p. 218]. Since $\Delta|z|^\alpha = \alpha^2|z|^{\alpha-2}$, we see that $\varphi_j(z) = |z|^\alpha$ satisfies the assumptions of Theorem 5.1 for all $\alpha \geq 4$ (so that φ is at least C^2).

(ii) Let $n \geq 2$ and consider the weight function

$$\varphi(z) = \sum_{j=1}^{n} |z_j|^{\alpha_j}, \tag{5.9}$$

where $\alpha_j \in \mathbb{R}$, $\alpha_j \geq 4$. Then $\varphi \in C^2(\mathbb{C}^n, \mathbb{R})$ and by the above

$$\lim_{z\to\infty} \operatorname{tr}(M_\varphi(z)) = \lim_{z\to\infty} \frac{1}{4} \sum_{j=1}^{n} \alpha_j^2 |z_j|^{\alpha_j - 2} = +\infty.$$

Therefore it follows from Theorem 5.1 that $N^\varphi_{0,q}$ with $0 \leq q \leq n-1$ is not compact while $N^\varphi_{0,n}$ is compact. Hence, the necessary conditions (3.1) and (4.3) fail to be sufficient for compactness of $N^\varphi_{0,q}$ for $0 \leq q \leq n-1$ in general. Of course, by Remark 4.2 and Theorem 5.1, the integral condition (4.3) is both necessary and sufficient for compactness of $N^\varphi_{0,n}$ for plurisubharmonic weights φ with $\operatorname{tr}(M_\varphi) \in A_\infty$, such as (5.9).

(iii) Using a variation of the above decoupled weights, one easily sees that, for $q > n/2$, there is a plurisubharmonic function $\varphi_q \colon \mathbb{C}^n \to \mathbb{R}$, such that $N^{\varphi_q}_{0,k}$ is compact precisely for $q \leq k \leq n$. Indeed, one may take

$$\varphi_q(z_1, \ldots, z_n) := |(z_1, \ldots, z_{q-1})|^4 + |(z_q, \ldots, z_n)|^4.$$

Then both of the spaces $A^2(\mathbb{C}^{q-1}, e^{-\varphi_1})$ and $A^2(\mathbb{C}^{n-q+1}, e^{-\varphi_2})$, where $\varphi_1 \colon \mathbb{C}^{q-1} \to \mathbb{R}$, $\varphi(z) := |z|^4$, and $\varphi_2 \colon \mathbb{C}^{n-q+1} \to \mathbb{R}$, $\varphi_2(z) := |z|^4$, have infinite dimension by the result of Shigekawa quoted in Remark 3.2. Moreover, the $\overline{\partial}$-Neumann operators $N^{\varphi_j}_{0,q}$ are compact for $q \geq 1$ and $j \in \{1, 2\}$, as is easily deduced by verifying (1.5). Since $n - q + 1 < n/2 + 1$ implies $n - q + 1 \leq n/2 \leq q - 1$, one obtains from [2, Theorem 5.2] that $N^\varphi_{0,k}$ is compact exactly for $k = q - 1 + j$ with $j \geq 1$, as claimed.

References

[1] N. Aronszajn. A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. *J. Math. Pures Appl. (9)*, 36:235–249, 1957. ISSN 0021-7824.

[2] F. Berger. Essential spectra of tensor product Hilbert complexes, and the $\overline{\partial}$-Neumann problem on product manifolds. *Journal of Functional Analysis*, 2016. doi: 10.1016/j.jfa.2016.06.004.

[3] J. Brüning and M. Lesch. Hilbert complexes. *J. Funct. Anal.*, 108(1):88–132, 1992. ISSN 0022-1236. doi: 10.1016/0022-1236(92)90147-B.

[4] D. Chakrabarti. Spectrum of the complex Laplacian on product domains. *Proc. Amer. Math. Soc.*, 138(9): 3187–3202, 2010. ISSN 0002-9939. doi: 10.1090/S0002-9939-10-10522-X.

[5] D. Chakrabarti and M.-C. Shaw. The Cauchy-Riemann equations on product domains. *Math. Ann.*, 349(4): 977–998, 2011. ISSN 0025-5831. doi: 10.1007/s00208-010-0547-x.

[6] G. M. Dall'Ara. *Matrix Schrödinger operators and weighted Bergman kernels.* PhD thesis, 2014.

[7] G. M. Dall'Ara. Coercivity of weighted Kohn Laplacians: the case of model monomial weights in $\mathbb{C}^2$, 2015.

[8] J.-P. Demailly. *Complex analytic and differential geometry.* 2012. Available online at https://www-fourier.ujf-grenoble.fr/~demailly/.

[9] M. P. Gaffney. Hilbert space methods in the theory of harmonic integrals. *Trans. Amer. Math. Soc.*, 78: 426–444, 1955. ISSN 0002-9947.

[10] F. Haslinger. Compactness for the $\overline{\partial}$-Neumann problem: a functional analysis approach. *Collect. Math.*, 62(2): 121–129, 2011. ISSN 0010-0757. doi: 10.1007/s13348-010-0013-9.

[11] F. Haslinger. *The $\overline{\partial}$-Neumann problem and Schrödinger operators*, volume 59 of *De Gruyter Expositions in Mathematics*. De Gruyter, Berlin, 2014. ISBN 978-3-11-031530-1; 978-3-11-031535-6. doi: 10.1515/9783110315356.

[12] F. Haslinger and B. Helffer. Compactness of the solution operator to $\overline{\partial}$ in weighted L^2-spaces. *J. Funct. Anal.*, 243(2):679–697, 2007. ISSN 0022-1236. doi: 10.1016/j.jfa.2006.09.004.

[13] L. Hörmander. L^2 estimates and existence theorems for the $\bar{\partial}$ operator. *Acta Math.*, 113:89–152, 1965. ISSN 0001-5962.

[14] D. Huybrechts. *Complex geometry*. Universitext. Springer-Verlag, Berlin, 2005. ISBN 3-540-21290-6. An introduction.

[15] A. Iwatsuka. Magnetic Schrödinger operators with compact resolvent. *J. Math. Kyoto Univ.*, 26(3):357–374, 1986. ISSN 0023-608X.

[16] X. Ma and G. Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007. ISBN 978-3-7643-8096-0.

[17] N. Marco, X. Massaneda, and J. Ortega-Cerdà. Interpolating and sampling sequences for entire functions. *Geom. Funct. Anal.*, 13(4):862–914, 2003. ISSN 1016-443X. doi: 10.1007/s00039-003-0434-7.

[18] J. Marzo and J. Ortega-Cerdà. Pointwise estimates for the Bergman kernel of the weighted Fock space. *J. Geom. Anal.*, 19(4):890–910, 2009. ISSN 1050-6926. doi: 10.1007/s12220-009-9083-x.

[19] G. Rozenblum and N. Shirokov. Infiniteness of zero modes for the Pauli operator with singular magnetic field. *J. Funct. Anal.*, 233(1):135–172, 2006. ISSN 0022-1236. doi: 10.1016/j.jfa.2005.08.001.

[20] J. Ruppenthal. Compactness of the $\overline{\partial}$-Neumann operator on singular complex spaces. *J. Funct. Anal.*, 260(11):3363–3403, 2011. ISSN 0022-1236. doi: 10.1016/j.jfa.2010.12.022.

[21] I. Shigekawa. Spectral properties of Schrödinger operators with magnetic fields for a spin $\frac{1}{2}$ particle. *J. Funct. Anal.*, 101(2):255–285, 1991. ISSN 0022-1236. doi: 10.1016/0022-1236(91)90158-2.

[22] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. ISBN 0-691-03216-5. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.

[23] E. J. Straube. *Lectures on the $\mathscr{L}^2$-Sobolev theory of the $\overline{\partial}$-Neumann problem*. ESI Lectures in Mathematics and Physics. European Mathematical Society (EMS), Zürich, 2010. ISBN 978-3-03719-076-0. doi: 10.4171/076.

[24] R. O. Wells, Jr. *Differential analysis on complex manifolds*. Prentice-Hall, Inc., Englewood Cliffs, N.J., 1973. Prentice-Hall Series in Modern Analysis.

Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, A-1090 Wien, Austria
E-mail address: franz.berger2@univie.ac.at, friedrich.haslinger@univie.ac.at